



## Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact [support@jstor.org](mailto:support@jstor.org).

(2). Let  $FG$  be the random line,  $AH=x$ ,  $DG=y$ .

Then  $FH=\frac{1}{3}\sqrt{3}x$ ,  $FG=\sqrt{[(a\sqrt{3}-x)^2+(y-\frac{1}{3}\sqrt{3}x)^2]}=l$ .

$$\Delta = \text{average length} = \frac{\int_0^{a\sqrt{3}} \int_0^a l dx dy}{\int_0^{a\sqrt{3}} \int_0^a dx dy}$$

$$\begin{aligned} \Delta &= \frac{1}{a^2\sqrt{3}} \int_0^{a\sqrt{3}} \int_0^a \sqrt{[(a\sqrt{3}-x)^2+(y-\frac{1}{3}\sqrt{3}x)^2]} dx dy \\ &= \frac{1}{12a^2\sqrt{3}} \int_0^{a\sqrt{3}} \left[ 4(a\sqrt{3}-x)^2 + x\sqrt{[9a^2+(4x-3\sqrt{3}a)^2]} \right. \\ &\quad \left. - 6(a\sqrt{3}-x)^2 \log \left( \frac{\sqrt{[9a^2+(4x-3\sqrt{3}a)^2]} - 2x}{6(\sqrt{3}-x)} \right) \right] dx. \\ \therefore \Delta &= (a/16)[6+2\sqrt{3}+4\log 3+\log(3+2\sqrt{3})]. \end{aligned}$$

82. Proposed by B. F. FINKEL, A. M., M. Sc., Professor of Mathematics and Physics, Drury College, Springfield, Mo.

Find the average area of the quadrilateral formed by joining the extremities of two chords perpendicular to each other and passing through a point at a distance  $a$  from the center of a circle radius  $R$ .

Solution by G. B. M. ZERR, A. M., Ph. D., Professor of Mathematics and Science, Chester High School, Chester, Pa.; and the PROPOSER.

Let  $\Delta$  = average area,  $\angle BPC = \theta$ ,  $P$  the given point,  $C$  the center of the circle,  $CP = a$ ,  $CB = R$ .

CASE I.  $a < R$ ,  $CG = a \sin \theta$ ,  $CF = a \cos \theta$ .

$\therefore AB = 2\sqrt{(R^2 - a^2 \sin^2 \theta)}$ ,  $DE = 2\sqrt{(R^2 - a^2 \cos^2 \theta)}$ .

Area  $ADBE = 2\sqrt{[(R^2 - a^2 \sin^2 \theta)(R^2 - a^2 \cos^2 \theta)]}$   
 $= 2R^2 \sqrt{[(1 - e^2 \sin^2 \theta)(1 - e^2 \cos^2 \theta)]}$  where  $e^2 = a^2/R^2$ .

$$\therefore \Delta = 2R^2 \int_0^{\frac{1}{2}\pi} \sqrt{[(1 - e^2 \sin^2 \theta)(1 - e^2 \cos^2 \theta)]} d\theta / \int_0^{\frac{1}{2}\pi} d\theta$$

$$= \frac{2R^2}{\pi} (2 - e^2) E\left(\frac{e^2}{2 - e^2}, \frac{1}{2}\pi\right).$$

(See page 356, No. 10, Vol. I, for above integration.)

CASE II.  $a > R$  and  $a < R\sqrt{2}$ ,  $PF = a \cos \theta$ ,  $PG = a \sin \theta$ .


$PB = a \cos \theta + \sqrt{(R^2 - a^2 \sin^2 \theta)}$ ,  $PA = a \cos \theta - \sqrt{(R^2 - a^2 \sin^2 \theta)}$ .

$PE = a \sin \theta + \sqrt{(R^2 - a^2 \sin^2 \theta)}$ ,  $PD = a \sin \theta - \sqrt{(R^2 - a^2 \cos^2 \theta)}$ .


Area  $ABED = \frac{1}{2}(BP \cdot PE - AP \cdot PD)$

$$= a[\sqrt{(R^2 - a^2 \sin^2 \theta)} \sin \theta + \sqrt{(R^2 - a^2 \cos^2 \theta)} \cos \theta].$$

The limits of  $\theta$  are  $\frac{1}{2}\pi - \sin^{-1}(R/a) = \theta''$  and  $\sin^{-1}(R/a) = \theta'$ .



$$\begin{aligned}
\therefore \Delta &= aR \int_{\theta'}^{\theta''} [\sqrt{1-e^2 \sin^2 \theta} \sin \theta + \sqrt{1-e^2 \cos^2 \theta} \cos \theta] d\theta / \int_{\theta'}^{\theta''} d\theta \\
&= \frac{aR}{4 \sin^{-1}(R/a) - \pi} \left[ \sin \theta' \sqrt{1-e^2 \cos^2 \theta'} \right. \\
&\quad - \sin \theta'' \sqrt{1-e^2 \cos^2 \theta''} - \cos \theta' \sqrt{1-e^2 \sin^2 \theta'} \\
&\quad \left. + \cos \theta'' \sqrt{1-e^2 \sin^2 \theta''} \right] \\
&+ \frac{1-e^2}{e} \log \left( \frac{e \sin \theta' + \sqrt{1-e^2 \cos^2 \theta'}}{e \sin \theta'' + \sqrt{1-e^2 \cos^2 \theta''}} \right) \\
&\quad - \frac{1-e^2}{e} \log \left( \frac{e \cos \theta' + \sqrt{1-e^2 \sin^2 \theta'}}{e \cos \theta'' + \sqrt{1-e^2 \sin^2 \theta''}} \right) \Big].
\end{aligned}$$



But  $\sin \theta' = \cos \theta'' = R/a$ ,  $\sin \theta'' = \cos \theta' = [\sqrt{a^2 - R^2}]/a$ .  
 $e^2 \sin^2 \theta' = e^2 \cos^2 \theta'' = 1$ ,  $e^2 \sin^2 \theta'' = e^2 \cos^2 \theta' = [(a^2 - R^2)/R^2] = e^2 - 1$ .  
 Whence by substitution and reduction we get

$$\Delta = \frac{R^2}{2 \sin^{-1}(R/a) - \frac{1}{2} \pi} \left[ \sqrt{1-e^2} + (1-e^2) \log \left( \frac{1 + \sqrt{1-e^2}}{1/(e^2-1)} \right) \right].$$

83. Proposed by F. P. MATZ, M. Sc., Ph. D., Professor of Mathematics and Astronomy, Irving College, Mechanicsburg, Pa.

Find the average area of all ellipses whose semi-axis major is  $a$ .

I. Solution by J. W. YOUNG, Fellow and Assistant, Ohio State University, Columbus, Ohio.

The area of an ellipse whose major-axis is  $a$ , and whose minor-axis is  $b$ , is  $\pi ab$ . We must find the average of all possible values of this expression as  $b$  varies from zero to  $a$ .

$$\begin{aligned}
&\frac{\pi a \int_0^a b db}{\int_0^a db} \\
\therefore \text{Average required} &= \frac{\pi a \int_0^a b db}{\int_0^a db} = \frac{1}{2} \pi a^2,
\end{aligned}$$

$= \frac{1}{2}$  the area of the circle whose radius is the major-axis of the ellipse.

II. Solution by G. B. M. ZERR, A. M., Ph. D., Professor of Mathematics and Science, Chester High School, Chester, Pa.

1. Let  $x$  = semi-conjugate axis. Then average area

$$\begin{aligned}
&\frac{\int_0^a x dx}{\int_0^a dx} \\
&= \frac{\pi a \int_0^a x dx}{\int_0^a dx} = \frac{1}{2} \pi a^2 = 1.5708 a^2.
\end{aligned}$$